



Mathematical Induction

Principle of Mathematical Induction

Does \mathbf{Z}^+ have any distinct property against \mathbf{Q}^+ and \mathbf{R}^+ ?

$$\mathbf{Z}^+ = \{ x \in \mathbf{Z} \mid x > 0 \} = \{ x \in \mathbf{Z} \mid x \geq 1 \}$$

$$\mathbf{Q}^+ = \{ x \in \mathbf{Q} \mid x > 0 \}, \quad \mathbf{R}^+ = \{ x \in \mathbf{R} \mid x > 0 \}$$

■ The well-ordering principle:

Every nonempty subset of \mathbf{N} contains a smallest element.

(\mathbf{N} is well-ordered.)

- Can be used to prove the principle of mathematical induction
- \mathbf{R}^+ is not well-ordered.

Principle of Mathematical Induction

- **Theorem:** The Principle of Mathematical Induction

Let $P(n)$ be a proposition for a natural number n .

- If $P(0)$ is true; and
- If $(\forall k \in \mathbf{N}) (P(k) \rightarrow P(k + 1))$ is true;

Then, $(\forall n \in \mathbf{N}) P(n)$ is true

- Consider applying the Modus Ponens,

$P(0)$

$P(0) \rightarrow P(1) \quad P(1)$

$P(1) \rightarrow P(2) \quad P(2)$

.....

...

$P(k) \rightarrow P(k + 1) \quad P(k + 1)$

Principle of Mathematical Induction

■ *Proof:*

Let $F = \{ t \in \mathbf{N} \mid P(t) \text{ is false} \}$.

Suppose $F \neq \emptyset$.

Then, there must be a smallest element $s \in F$ by the well-ordering principle.

Since $P(0)$ is true, $s \neq 0$.

So, $s > 0$ and thus $s - 1 \in \mathbf{N}$.

With $s - 1 \notin F$ we have $P(s - 1)$ true.

Therefore, $P((s - 1) + 1) = P(s)$ is true, which is a contradiction.

Examples

- For all $n \in \mathbf{Z}^+$,

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

- *Proof*:

(Basis step) For $n = 1$

LHS = 1, RHS = 1. So, LHS = RHS.

(Inductive step)

We want to show that

$$\sum_{i=1}^n i = \frac{n(n+1)}{2} \Rightarrow \sum_{i=1}^{n+1} i = \frac{(n+1)(n+2)}{2}$$

Examples

- *Proof:*

Assume that $\sum_{i=1}^n i = \frac{n(n+1)}{2}$ (Inductive Hypothesis, AP)

Then,

$$\begin{aligned}\sum_{i=1}^{n+1} i &= (n+1) + \sum_{i=1}^n i \\ &= (n+1) + \frac{n(n+1)}{2} \quad \text{(by inductive hypothesis)}\end{aligned}$$

$$= \frac{2(n+1) + n(n+1)}{2}$$

$$= \frac{(n+1)(n+2)}{2}$$

(Now, we want to apply the CP rule.)

Examples

- Let $r \neq 0$ and $r \neq 1$

$$\sum_{i=0}^n r^i = \frac{r^{n+1} - 1}{r - 1}$$

- *Proof:*

(Basis step) For $n = 0$

LHS = 1, RHS = 1. So, LHS = RHS.

(Inductive step)

Assume that $\sum_{i=0}^n r^i = \frac{r^{n+1} - 1}{r - 1}$ (Additional premise, i.e., I.H.)

Examples

- *Proof:*

We want to show that $\sum_{i=0}^{n+1} r^i = \frac{r^{n+2} - 1}{r - 1}$

and we will use the CP rule.

$$\begin{aligned}\sum_{i=0}^{n+1} r^i &= r^{n+1} + \sum_{i=0}^n r^i \\ &= r^{n+1} + \frac{r^{n+1} - 1}{r - 1} \quad (\text{by I.H.}) \\ &= \frac{r^{n+2} - r^{n+1} + r^{n+1} - 1}{r - 1} = \frac{r^{n+2} - 1}{r - 1}\end{aligned}$$

Examples

- For every $n \in \mathbf{N}$, $7^n - 2^n$ is divisible by 5.

- *Proof*:

(Basis step) For $n = 0$

$7^0 - 2^0 = 0$ is divisible by 5.

(Inductive step)

Let $7^n - 2^n$ be divisible by 5.

Then,

$$\begin{aligned}7^{n+1} - 2^{n+1} &= 7 \cdot (7^n - 2^n) + 7 \cdot 2^n - 2^{n+1} \\ &= 7 \cdot (7^n - 2^n) + 2^n \cdot (7 - 2)\end{aligned}$$

Examples

- *Proof:*

Since $(7^n - 2^n)$ is divisible by 5 by the inductive hypothesis,
 $7 \cdot (7^n - 2^n)$ is divisible by 5.

Also, $2^n \cdot (7 - 2)$ is divisible by 5.

Therefore, $7^{n+1} - 2^{n+1}$ is divisible by 5.

Examples

■ If S is a finite set then $|\wp(S)| = 2^{|S|}$.

■ *Proof:*

(Basis step) For $S = \emptyset$

$$\text{LHS} = |\wp(\emptyset)| = |\{\emptyset\}| = 1 = 2^0 = 2^{|\emptyset|} = \text{RHS}.$$

(Inductive step)

Let $|\wp(S)| = 2^{|S|} = 2^n$ for $S = \{a_1, a_2, \dots, a_n\}$.

We want to prove that $|\wp(S')| = 2^{|S'|} = 2^{n+1}$

where $S' = \{a_1, a_2, \dots, a_n, a_{n+1}\}$.

Examples

■ *Proof:*

We know that if $X \subseteq S$ then $X \subseteq S'$, which means that every subset of S is a subset of S' .

But, note that $X \cup \{a_{n+1}\} \subseteq S'$ for any $X \subseteq S$ and there is no other subset of S' in addition to these subsets.

Therefore, the number of subsets of S' is twice that of S , i.e.,

$$|\wp(S')| = 2 \cdot |\wp(S)| = 2 \cdot 2^{|S|} = 2 \cdot 2^n = 2^{n+1} = 2^{|S'|}.$$

Examples

- The number of left parenthesis is equal to the number of right parenthesis in a propositional well-formed formula.

- *Proof:*

Let $\#L(\mathbf{F})$ and $\#R(\mathbf{F})$ denote the number of left parenthesis and the number of right parenthesis of a wff \mathbf{F} .

(Basis Step)

Since any propositional variable or constant S has no parenthesis by the basis clause of the inductive definition of a wff, $\#L(S) = \#R(S)$.

Examples

- *Proof:*

(Inductive Step)

Let P and Q be two wffs such that

$$\#L(P) = \#R(P) \text{ and } \#L(Q) = \#R(Q).$$

Let \mathbf{F} be any one of the formulas defined by the inductive clause of the inductive definition of a wff, that is, $(\neg P)$, $(P \vee Q)$, $(P \wedge Q)$, $(P \rightarrow Q)$, and $(P \leftrightarrow Q)$.

If $\mathbf{F} = (\neg P)$, then $\#L(\mathbf{F}) = \#L(P) + 1$ and $\#R(\mathbf{F}) = \#R(P) + 1$.

Therefore, $\#L(\mathbf{F}) = \#R(\mathbf{F})$.

Examples

- *Proof:*

On the other hand, if \mathbf{F} is $(P \vee Q)$, $(P \wedge Q)$, $(P \rightarrow Q)$, or $(P \leftrightarrow Q)$, then

$$\#L(\mathbf{F}) = \#L(P) + \#L(Q) + 1 \text{ and}$$

$$\#R(\mathbf{F}) = \#R(P) + \#R(Q) + 1.$$

Again, since $\#L(P) = \#R(P)$ and $\#L(Q) = \#R(Q)$,

$$\#L(\mathbf{F}) = \#R(\mathbf{F}).$$